

Q17

Is depletion of the ozone layer the principal cause of global climate change?

No, ozone depletion is not the principal cause of global climate change. Ozone depletion and global climate change are linked because both ozone-depleting substances and their substitutes are greenhouse gases. The best estimate is that stratospheric ozone depletion has led to a small amount of surface cooling. Conversely, increases in tropospheric ozone and other greenhouse gases lead to surface warming. The effect on global climate change from stratospheric ozone depletion is small compared to the warming from the greenhouse gases responsible for observed global climate change. Since the early 1980s, the Antarctic ozone hole has contributed to changes in Southern Hemisphere surface climate through effects on the atmospheric circulation.

While stratospheric ozone depletion is not the principal cause of climate change, aspects of ozone depletion and climate change are closely linked. Both processes involve gases released to the atmosphere by human activities. The links are best understood by examining the contribution to climate change of the gases involved: ozone; ozone-depleting substances (or halogen source gases) and their substitutes; and other leading greenhouse gases.

Greenhouse gases and the radiative forcing of climate.

The warming of Earth's surface and troposphere by the Sun is enhanced by the presence of greenhouse gases (GHGs). The natural abundances of GHGs in Earth's atmosphere absorb outgoing infrared radiation, trapping heat in the atmosphere and warming the surface. The most important natural GHG is water vapor. Without this natural greenhouse effect, Earth's surface would be much colder than current conditions. Human activities have led to significant increases in the atmospheric abundances of a number of long-lived and short-lived GHGs since 1750, the start of the Industrial Era, leading to warming of Earth's surface and associated climate changes. This group includes carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O), tropospheric ozone, and halocarbons. Ozone-depleting substances (ODSs) and their substitutes make up a large fraction of the halocarbons in today's atmosphere. Increases in the abundances of these gases from human activities cause more outgoing infrared radiation to be absorbed and reemitted back to the surface, further warming the atmosphere and surface. This change in Earth's energy balance caused by human activities is called a *radiative forcing of climate* or, more simply, a *climate forcing*. The magnitude of this energy imbalance is usually evaluated at the top of the troposphere (tropopause) and is expressed using units of watts per square meter (W/m²). The potential for climate change rises as this radiative forcing increases.

A summary of radiative forcings of climate in 2019 resulting from the increases in the principal long-lived and short-lived GHGs from human activities since 1750 is shown in **Figure Q17-1**. Positive forcings lead to warming and negative forcings lead to cooling of Earth's surface. Climate forcings also lead to other changes, for example reductions in glacier and sea-ice extent, variations in precipitation patterns, and more extreme weather events. International climate assessments conclude that much of the observed

surface warming and changes in other climate parameters over the last several decades are due to increases in the atmospheric abundances of CO₂ and other GHGs, which result from a variety of human activities.

Carbon dioxide, methane, and nitrous oxide. All three of these GHGs have both human and natural sources. The accumulation of CO₂ since 1750 represents the largest climate forcing caused by human activities. Carbon dioxide concentrations continue to increase in the atmosphere primarily as the result of burning fossil fuels (coal, oil, and natural gas) for energy and transportation, as well as from cement manufacturing. The global mean atmospheric abundance of CO₂ exceeded 416 parts per million (ppm) in 2021, which is about 50% larger than the abundance of CO₂ present in 1750. Carbon dioxide is considered a long-lived gas, since a significant fraction remains in the atmosphere 100 to 1000 years after emission.

Methane is a *short-lived* climate gas (atmospheric lifetime of about 12 years). Sources related to human activities include livestock, fossil fuel extraction and use, rice agriculture, and landfills. Natural sources include wetlands, termites, and oceans. The global mean atmospheric abundance of CH₄ has more than doubled since 1750.

Nitrous oxide is a *long-lived* climate gas (atmospheric lifetime of about 109 years). The largest source related to human activities is agriculture, especially the use of fertilizer. Microbial processes in soils that are part of natural biogeochemical cycles represent the largest natural source. In the stratosphere, nitrous oxide is the principal source of reactive nitrogen species that participate in ozone destruction cycles (see Q8). The global mean atmospheric abundance of nitrous oxide has increased by about 22% since 1750.

Halocarbons. Halocarbons in the atmosphere contribute to both ozone depletion and climate change. The halocarbons considered in Figures Q17-1 and **Q17-2** are gases containing chlorine, bromine, or fluorine atoms that are either controlled under the Montreal Protocol or are GHGs that fall under the auspices of the United Nations Framework Convention on Climate Change (UNFCCC). Historically, ODSs were the only halocarbons controlled under the Montreal Protocol. In 2016, the Kigali Amend-

Greenhouse Gases and Climate Change

Climate forcing due to changes in greenhouse gases caused by human activities between 1750 and 2019




Figure Q17-1. Radiative forcing of greenhouse gases and ozone depletion. Human activities since the start of the Industrial Era (around 1750) have caused increases in the atmospheric abundance of greenhouse gases (GHGs). Rising levels of GHGs lead to an increase in the radiative forcing (RF) of climate by trapping infrared radiation released by Earth's surface. Here, values of RF are for the time period from 1750 to 2019 and are expressed in units of watts per square meter (W/m^2); black whiskers on each bar show uncertainties. Positive values of RF (shown in red) contribute to global warming and negative values (shown in blue) contribute to climate cooling. The largest positive RFs are due to carbon dioxide (CO₂), methane (CH₄), tropospheric ozone (O₃), halocarbons, and nitrous oxide (N₂O). Halocarbons include all ozone-depleting substances, hydrofluorocarbons, and a few other gases (see Figure Q17-2). The RF due to ozone is shown as the separate response to changes in ozone within two layers of the atmosphere: the troposphere and stratosphere. Tropospheric ozone increases result from the emission of air pollutants and lead to a positive RF whereas stratospheric ozone depletion leads to a small RF that could be negative or positive.

ment to the Montreal Protocol established controls on the future production and consumption of certain hydrofluorocarbon (HFC) gases. Perfluorocarbons (PFCs) and sulfur hexafluoride (SF₆) are in the UNFCCC group of GHGs that now fall under the Paris Agreement. Perfluorocarbons are compounds that contain only carbon and fluorine atoms, such as carbon tetrafluoride (CF₄) and perfluoroethane (C₂F₆). Technically, SF₆ is not a halocarbon since it lacks carbon. However, the environmental effects of SF₆ are commonly examined with those of halocarbon gases since all of these compounds contain at least one halogen atom.

In 2019, the halocarbon contribution to the radiative forcing (RF) of climate was 0.41 W/m^2 , which is the fourth largest GHG forcing following carbon dioxide, methane, and tropospheric ozone (see Figure Q17-1). The contributions of individual halocarbon gases controlled by the Montreal Protocol are highlighted in Figure Q17-2. Within the halocarbons, CFC-12, CFC-11, and CFC-113 combined contribute the largest percentage (67%) to radiative forcing in 2019. The intermediate-term ODS substitutes, hydrochlorofluorocarbons (HCFCs), make the next largest contribution (15%). The long-term ODS substitutes HFCs plus PFCs and SF₆ contribute 13%. Finally, CCl₄ and CH₃CCl₃ contribute an additional 3% and the minor CFCs and halons contribute 2% to radiative forcing in 2019.

The large contribution of the CFCs to radiative forcing has been gradually decreasing following the decline in their atmospheric abundance and is expected to further decrease (see Figure Q15-1). Based on their long lifetimes, CFCs will still make a significant contribution, and most likely the largest contribution from ODSs, to the radiative forcing by halocarbons at the end of this century. Even with adherence to the provisions of the Kigali Amendment to the Montreal Protocol, the radiative forcing from HFCs is projected to increase for another two to three decades before starting to slowly decline (see Figure Q19-2).

Stratospheric and tropospheric ozone. Ozone in both the stratosphere and the troposphere absorbs infrared radiation emitted from Earth's surface, trapping heat in the atmosphere. Ozone also significantly absorbs solar ultraviolet (UV) radiation. As a result, increases or decreases in stratospheric or tropospheric ozone induce a climate forcing and, therefore, represent direct links between ozone and climate. Air pollution from a variety of human activities has led to increases in global tropospheric ozone (see Q2), causing a positive radiative forcing (warming) estimated to be +0.47 W/m^2 over the 1750–2019 time period, with a range of uncertainty spanning +0.24 to +0.70 W/m^2 (see Figure Q17-1). The large uncertainty in the climate forcing due to release of air pollutants reflects our limited knowledge of changes in the abundance of tropospheric ozone between 1750 and the mid-1950s as well

Radiative Forcing of Climate by Halocarbons

Climate forcing due to increases in controlled gases containing chlorine, bromine, and fluorine from human activities between 1750 and 2019




Figure Q17-2. Halocarbons and radiative forcing of climate. Halocarbon gases in the atmosphere represent an important contribution to the radiative forcing (RF) of climate since the start of the Industrial Era (see Figure Q17-1). Halocarbons are gases containing chlorine, bromine, or fluorine atoms, with at least one carbon atom, that contribute to RF by trapping infrared radiation released by Earth’s surface. The rise in RF between 1750 and 2019 is shown for all halocarbons controlled either under the Montreal Protocol (red) or included in the Paris Agreement (blue) along with the RF due to the rise in SF₆. Note that while SF₆ is technically not a halocarbon because it lacks any carbon atoms, it is an important halogen-containing gas in the atmosphere. Separate contributions to RF of each gas or group of gases are based upon atmospheric abundance histories and the radiative efficiency specific to each compound. The gases listed in the right-hand labels begin with the largest contribution in each group and proceed in descending order, except for the entry for minor CFCs and halons, which are shown as one total value. The individual RF terms add together to form the bottom bar, representing the total RF due to controlled halocarbons, PFCs, and SF₆. The RFs of CFC-11 and CFC-12, the largest halocarbon contributors, are decreasing and will continue to decline as CFCs are gradually removed from the atmosphere (see Figure Q15-1). In contrast, the total RF of HCFCs, the intermediate-term ODS substitute gases, is projected to grow for about another one to two decades before decreasing. HFCs are the long-term ODS substitute gases. With the October 2016 Kigali Amendment, the Montreal Protocol now controls future production and consumption of HFCs with high Global Warming Potentials. As a result, nearly all of the RF due to halogen-containing GHGs is now controlled by the Montreal Protocol (bottom bar). The future RF of climate due to HFCs is expected to peak in about two decades under the provisions of the Kigali Amendment (see Q19).

as the difficulty in modeling the complex chemical processes that control the production of tropospheric ozone.

On the other hand, rising abundances of ODSs in the atmosphere since the middle of the 20th century have led to decreases in stratospheric ozone, causing a negative radiative forcing of -0.02 W/m^2 (cooling) over the 1750–2019 time period, with a range of uncertainty spanning -0.15 to $+0.11 \text{ W/m}^2$ (see Figure Q17-1). The sign of the radiative forcing due to stratospheric ozone depletion is uncertain because this quantity is the difference between two terms of comparable magnitude, each of which has an associated uncertainty. The first term represents the trapping by ozone of outgoing infrared radiation released by the surface and lower atmosphere: this is a cooling term because less ozone results in less trapping of heat. The second term represents the absorption of solar UV radiation by ozone: this is a warming term because less ozone results in greater penetration of solar UV radiation into the lower atmo-

sphere (troposphere). This radiative forcing due to stratospheric ozone depletion will diminish in the coming decades, as ODSs are gradually removed from the atmosphere.

It is clear that stratospheric ozone depletion is not a principal cause of present-day global warming. First, the climate forcing from ozone depletion is small. Second, the total radiative forcing of climate from other GHGs such as carbon dioxide, methane, halocarbons, and nitrous oxide is large and positive, leading to warming (see Figure Q17-1). The total forcing from these other GHGs is the principal cause of the observed warming of Earth’s surface.

Ozone Depletion Potentials and Global Warming Potentials. A useful way of comparing the influence of individual emissions of halocarbons on ozone depletion and climate change is to compare Ozone Depletion Potentials (ODPs) and Global Warming Potentials (GWPs). The ODP and GWP are the effectiveness of an emission of a gas in causing ozone depletion and climate forcing,

Evaluation of Selected Ozone-Depleting Substances and Substitute Gases

Relative importance of equal mass emissions for ozone depletion and climate change




Figure Q17-3. ODPs and GWPs. The environmental impacts of ozone-depleting substances and their substitutes are commonly compared based upon their Ozone Depletion Potentials (ODPs) and Global Warming Potentials (GWPs) (see Table Q6-1). The ODPs and GWPs represent the magnitude of ozone depletion and climate forcing, respectively, of a given mass of gas emitted to the atmosphere, relative to that of CFC-11 (for ODP) or CO₂ (for GWP). Therefore, the ODP of CFC-11 and the GWP of CO₂ are assigned reference values of 1. The GWPs shown here are evaluated for a 100-year time interval after emission. The CFCs, halons, and HCFCs are ozone-depleting substances (ODSs) since they contain either chlorine or bromine (see Q6). HFCs, used as ODS substitutes, do not destroy ozone (ODPs equal zero) since they are mixtures of only hydrogen, fluorine, and carbon atoms. The ODPs of halons far exceed those of the CFCs, since all halons contain bromine. The GWPs of these gases span a wide range of values, from less than 1 (HFO-1234yf) to 14,700 (HFC-23).

respectively, relative to a reference gas (see Table Q6-1). The principal halocarbon gases are contrasted with each other in **Figure Q17-3**. The ODP of CFC-11 and the GWP of carbon dioxide are assigned reference values of 1. The CFCs and carbon tetrachloride all have ODPs near 1, indicating comparable effectiveness in causing ozone depletion per mass emitted. The principal halons have ODPs greater than 7, making them the most effective ozone-depleting substances per mass emitted. All HFCs have ODPs of zero since they contain no chlorine and bromine, and therefore do not directly cause ozone depletion (see Q6).

All halocarbons have non-zero GWPs and, therefore, contribute to the RF of climate. The GWP does not correspond strongly with

the ODP of a gas because these quantities depend on different chemical and physical properties of the molecule. For example, while HFC-143a does not destroy ozone (ODP equals zero), each kilogram emitted is about 6000 times more effective than a kilogram of carbon dioxide in causing climate forcing. When HFCs are released to the atmosphere, their contribution to climate forcing depends on their GWPs, which vary over a wide range (less than 1 to 15,000).

Montreal Protocol regulations have led to reductions in CFC emissions and increases in HCFC emissions (see Q15). As a result of these actions, the total RF from ODSs stopped increasing and is now slowly decreasing because HCFCs have lower GWPs than

CFCs (see Q18). However, the overall RF of all halocarbons is slowly increasing because of growing contributions from non-ODS gases (HFCs, PFCs, and SF₆). The growth in the HFC contribution will be limited by the provisions of the 2016 Kigali Amendment (see Q19). It is important to note that despite having a GWP that is small in comparison to many other halocarbons and other greenhouse gases, carbon dioxide is the most important greenhouse gas produced by human activities because its emissions are large, its atmospheric lifetime is long, and its atmospheric abundance

is far greater than those of all other greenhouse gases associated with human activities.

The Antarctic ozone hole and Southern Hemisphere climate. While stratospheric ozone depletion is not the principal cause of global climate change, the reoccurring Antarctic ozone hole has contributed to observed changes in climate parameters in the atmosphere and oceans of the Southern Hemisphere since the early 1980s. These research findings are explained in more detail in the box below.

The Antarctic Ozone Hole and Southern Hemisphere Surface Climate

Links between stratospheric ozone depletion and changes in surface climate were first found in research studies in the early 2000s, based on both observations and models. While increasing greenhouse gases (such as carbon dioxide, methane, and nitrous oxide) are the primary drivers of global climate change, the Antarctic ozone hole, which has occurred every spring since the early 1980s, was shown to contribute to observed changes in Southern Hemisphere (SH) surface climate during summer due to the effect of the ozone hole on atmospheric circulation.

The severe springtime depletion of ozone over the Antarctic leads to a strong cooling of the polar lower stratosphere persisting into early summer in the SH. This cooling increases the temperature contrast between the tropics and the polar region and strengthens stratospheric winds. As a result, in the SH there has been a poleward shift of tropospheric circulation features including the tropical Hadley cell (which determines the location of the subtropical dry zones) and the midlatitude jet stream (which is associated with weather systems). There is evidence from both models and observations that subtropical and midlatitude summer precipitation patterns in the SH have been affected by these changes. Model studies indicate that even though long-lived greenhouse gases that cause climate change exacerbate this shift in the summertime tropospheric circulation in the Southern Hemisphere, ozone depletion has been the dominant contributor to the observed changes since the early 1980s. Paleoclimate reconstructions suggest the current state of these climate features caused by the annually recurring ozone hole is unprecedented over the past 600 years.

Initial signs of the recovery of the ozone hole, particularly during September, are emerging (see Q10). During the 21st century, as the ozone hole further recovers due to the decline of stratospheric halogens, the ozone-depletion related climate impacts discussed above will lessen (see Q20). The influence of the ozone hole on SH circulation trends during summer has recently stabilized, due to the initial recovery of the area and depth of the ozone hole (see Q10). The Southern Hemisphere surface climate response to ozone depletion in other seasons is weaker than the summer response.